

Supplementary data

Title: Central adrenal insufficiency is rare in adults with Prader-Willi Syndrome

Authors: Anna G.W. Rosenberg, Karlijn Pellikaan, Christine Poitou, Anthony P. Goldstone, Charlotte Höybye, Tania Markovic, Graziano Grugni, Antonino Crinò, Assumpta Caixàs, Muriel Coupaye, Sjoerd A.A. van den Berg, Aart Jan van der Lely, Laura C.G. de Graaff

Corresponding author: Dr. Laura de Graaff, Internal Medicine, division of Endocrinology, Erasmus MC, University Medical Center Rotterdam, The Netherlands.
E-mail: l.degraaff@erasmusmc.nl. Telephone: +31-618843010

TABLE OF CONTENTS

Page 3 – 4	Table S1: Results of the multiple-dose metyrapone test
Page 5 – 6	Table S2: Results of the insulin tolerance test
Page 7	Table S3: Overview of adrenal function tests used to diagnose central adrenal insufficiency
Page 8	Figure S1: Relation between cortisol level (nmol/L) at baseline and increase in ACTH (pmol/L) during multiple-dose metyrapone test in patients with Prader-Willi syndrome
Page 9	Figure S2: ACTH levels (pmol/L) in Dutch patients with Prader-Willi syndrome who underwent the multiple-dose metyrapone test or insulin tolerance test
Page 10	References

Table S1. Results of the multiple-dose metyrapone test

Patient	Gender (M/F) ^a	Age (years)	BMI (kg/m ²) ^b	GH (Y/N) ^c	OAC/testosterone (Y/N) ^d	Baseline cortisol (nmol/L)	11-deoxycortisol (nmol/L) ^e	Delta ACTH (pmol/L) ^f
1	M	51.2	30.4	N	Y	235	407.9	6.1
2	M	37.8	48.1	N	Y	175	291.2	25.2
3	M	32.8	29.1	N	Y	350	520.2	41.3
4	M	20.1	41.9	N	Y	238	343.0	25.9
5	M	18.7	32.4	Y	Y	135	400.3	31.9
6	M	18.1	27.4	Y	Y	348	353.9	39.8
7	M	55.5	20.7	N	N	397	575.1	19.4
8	M	40.9	26.9	N	N	480	349.6	12.7
9	M	33.5	32.0	N	N	292	397.7	49.1
10	M	21.9	44.4	N	Y	252	279.9	21.9
11	M	27.6	26.3	N	Y	320	550.8	82.9
12	M	50.1	25.7	N	Y	395	390.9	15.8
13	M	42.9	25.0	N	N	301	355.0	10.9
14	M	35.7	24.6	N	Y	337	573.9	70.3
15	M	34.4	38.1	N	Y	428	358.3	43.8
16	M	29.2	27.8	N	Y	284	379.9	61.1
17	M	22.2	26.2	Y	Y	339	484.2	25.9
18	M	21.3	20.1	Y	Y	253	285.4	28.3
19	M	21.8	26.0	Y	Y	274	483.5	34.9
20	M	19.6	20.0	Y	Y	395	363.0	70.7
21	M	18.7	28.4	Y	Y	208	451.2	40.1
22	F	25.2	31.0	N	Y	709	660.7	13.2
23	F	21.8	34.4	N	N	126	495.3	40.2
24	F	20.9	36.9	Y	Y	508	425.3	13.7
25	F	29.9	37.7	N	Y	579	499.1	9.5

Patient	Gender (M/F)	Age (years)	BMI (kg/m ²)	GH (Y/N)	OAC/testosterone (Y/N)	Baseline cortisol (nmol/L)	11-deoxycortisol (nmol/L)	Delta ACTH (pmol/L)
26	F	29.7	26.8	Y	N	295	454.4	118.9
27	F	19.1	31.5	Y	Y	480	298.2	31.0
28	F	39.0	32.3	N	Y	233	544.3	73.6
29	F	29.1	32.8	Y	Y	345	500.3	71.0
30	F	27.6	33.9	Y	Y	167	476.2	4.1
31	F	22.5	21.7	Y	Y	601	487.9	44.0
32	F	22.3	27.2	N	Y	317	429.0	27.6
33	F	19.9	21.2	Y	Y	477	600.0	43.7
34	F	18.9	28.5	Y	Y	764	547.3	98.5
35	F	21.2	26.6	Y	Y	245	510.9	30.5
36	F	25.4	23.9	Y	Y	384	471.0	12.0
37	M	28.0	34.3	N	Y	306	391.2	-1.4
38	F	38.5	34.1	Y	N	216	247.8	20.5
39	F	29.5	31.5	N	Y	202	567.1	58.9
40	M	37.1	57.0	N	N	414	381.1	12.2
41	F	20.5	49.7	N	Y	508	569.4	91.9
42	M	21.4	20.6	Y	Y	273	401.7	28.2
43	M	51.1	29.4	N	N	350	334.0	39.9
44	M	19.4	24.7	Y	Y	331	694.0	61.5
45	M	22.9	25.6	Y	N	356	386.7	46.9
46	F	18.2	25.6	N	N	239	603.1	106.9

Y/N: yes/no. ^a M: male; F: female. ^b BMI: body mass index (kg/m²). ^c GH: current growth hormone treatment. ^d OAC: current use of oral estrogen or progesterone; testosterone: current use of testosterone. ^e 11-deoxycortisol (nmol/L) during multiple-dose metyrapone test. Cut-off for central adrenal insufficiency is <230 nmol/L. ^f Increase in ACTH (pmol/L) level during multiple-dose metyrapone test.

Table S2. Results of the insulin tolerance test

Patient	Gender (M/F) ^a	Age (years)	BMI (kg/m ²) ^b	GH (Y/N) ^c	OAC/testosterone (Y/N) ^d	Baseline cortisol (nmol/L)	Peak cortisol (nmol/L) ^e	Glucose (mmol/L)	Delta ACTH (pmol/L) ^f
Dutch patients									
47	M	22.4	28.3	N	N	254	662	2.4	35.5
48	M	37.8	28.0	N	Y	306	734	1.9	28.1
49	M	32.1	21.8	N	Y	242	828	1.7	52.2
50	M	30.7	24.3	N	Y	153	552	1.4	90.5
51	M	26.3	28.7	N	N	165	687	1.7	44.4
52	M	25.7	29.1	Y	Y	119	532	1.9	63.4
53	F	47.3	40.1	N	Y	450	883	1.5	64.1
54	F	31.8	34.6	Y	N	133	778	2.1	61.5
55	F	55.3	21.8	N	N	502	717	1.9	21.2
56	M	34.8	28.0	N	Y	306	734	1.9	90.5
Swedish patients									
57	M	20.0	21.2	N	Y	177	632	2.6	NA
58	M	22.0	44.4	N	Y	181	822	1.6	NA
59	M	25.0	24.2	N	Y	194	502	1.7	NA
60	M	27.0	27.5	N	Y	265	703	1.9	NA
61	F	31.0	36.5	N	N	190	742	1.4	NA
62	M	36.0	32.7	N	Y	175	771	1.2	NA
French patients									
63	M	18.0	47.8	N	NA	229	665	1.6	NA
64	M	20.0	49.7	N	NA	235	563	2.0	NA
65	M	20.0	28.3	Y	NA	378	494	2.0	NA

Patient	Gender (M/F)	Age (years)	BMI (kg/m ²)	GH (Y/N)	OAC/testosterone (Y/N)	Baseline cortisol (nmol/L)	Peak cortisol (nmol/L)	Glucose (mmol/L)	Delta ACTH (pmol/L)
66	F	29.0	32.0	N	NA	229	944	1.5	NA
67	F	24.0	31.8	N	NA	102	712	1.6	NA
68	F	28.0	58.2	N	NA	166	759	1.7	NA
69	F	38.0	27.3	N	NA	384	1021	0.9	NA
70	F	23.0	20.3	N	NA	116	789	2.2	NA
71	F	19.0	43.6	N	NA	132	601	1.4	NA
72	F	18.0	24.0	N	NA	240	869	0.6	NA
British patients									
73	F	18.3	33.8	N	Y	327	817	2.1	NA
74	M	30.9	24.1	N	N	242	603	1.1	NA
75	F	19.3	26.9	Y	Y	166	535	1.4	NA
76	F	19.4	29.9	N	N	179	510	1.0	NA
77	F	20.6	30.7	N	Y	148	455	1.7	NA
78	M	23.7	34.9	Y	N	296	538	1.7	NA
79	M	19.3	24.8	N	N	93	502	1.5	NA
80	F	19.2	35.2	N	Y	545	971	1.8	NA
81	M	26.1	62.0	N	N	119	467	1.5	NA
82	M	24.7	57.2	N	N	120	496	1.4	NA


Y/N: yes/no. NA: Not available. ^a M: male; F: female. ^b BMI: body mass index (kg/m²). ^c GH: current growth hormone treatment. ^d OAC: current use of oral estrogen or progesterone; testosterone: current use of testosterone. ^e Peak cortisol (nmol/L) level during insulin tolerance test. Cut-off for central adrenal insufficiency is <500 nmol/L (Dutch, French and Swedish patients) or <450 nmol/L (British patients). ^f Increase in ACTH (pmol/L) level during insulin tolerance test.

Table S3. Overview of adrenal function tests used to diagnose central adrenal insufficiency

Test	Test principle	Sensitivity (%)	Specificity (%)	Disadvantage
LDSST^{a 1}	Synthetic ACTH administration induces release of cortisol from the adrenal gland.	57	95	<ul style="list-style-type: none"> Adequate for the diagnosis of primary adrenal insufficiency, but lack of consensus about use for the diagnosis of <i>central</i> adrenal insufficiency.
HDSST^{b 1}		61	95	
MTP^{c 2}	Metirapone decreases cortisol production by inhibiting the 11 β -hydroxylase steroid enzyme. This stimulates ACTH production, leading to increased 11-deoxycortisol levels.	88-100	88-100	<ul style="list-style-type: none"> Risk of acute adrenal insufficiency precipitated by metirapone ingestion Should be conducted on an inpatient basis.
ITT^d	Insulin induces hypoglycaemia, which causes a release in ACTH and growth hormone (GH), leading to increased levels of cortisol.	Golden standard	Golden standard	<ul style="list-style-type: none"> Risks associated with hypoglycaemia Intravenous access at two different sites needed, which is impossible in many patients with PWS Labour intensive, time consuming, unpleasant for patient


^a LDSST: Low-dose short synacthen test. ^b HDSST: High-dose short synacthen test. ^c MTP: Metirapone test. ^d ITT: Insulin tolerance test.

Figure S1. Relation between cortisol level (nmol/L) at baseline and increase in ACTH (pmol/L) during multiple-dose metyrapone test in patients with Prader-Willi syndrome



Relationship between baseline cortisol (nmol/L) and increase in ACTH (pmol/L) in Dutch patients with Prader-Willi syndrome who underwent multiple-dose metyrapone test. N = 46. Spearman's rho was 0.01 ($P = 0.97$).

Figure S2. ACTH levels (pmol/L) in Dutch patients with Prader-Willi syndrome who underwent the multiple-dose metyrapone test or insulin tolerance test



Spaghetti plot of the ACTH levels (pmol/L) in patients with Prader-Willi syndrome who underwent the multiple-dose metyrapone test (MTP; black line) or insulin tolerance test (ITT; blue line). For MTP test, $n = 46$. For ITT, $n = 26$. Median (range) ACTH level before and after metyrapone administration (MTP) were 3.5 (1.3 – 16.2) pmol/L and 37.7 (2.8 – 132.0) pmol/L. Median (range) ACTH level before and after insulin administration (ITT) were 3.3 (1.1 – 6.2) pmol/L and 61.2 (23.8 – 93.5) pmol/L. The grey arrow represents the ACTH cut-off value of 33 pmol/L, used for interpretation of the MTP in the Dutch pediatric study.⁴ This would classify 21 patients with sufficient increase in 11-deoxycortisol levels as ‘adrenal insufficient’.

References supplementary data

1. Dorin RI, Qualls CR, Crapo LM. Diagnosis of adrenal insufficiency. *Ann Intern Med* 2003;139:194-204.
2. Fiad TM, Kirby JM, Cunningham SK, McKenna TJ. The overnight single-dose metyrapone test is a simple and reliable index of the hypothalamic-pituitary-adrenal axis. *Clin Endocrinol (Oxf)* 1994;40:603-9.
3. Gibney J, Healy ML, Smith TP, McKenna TJ. A simple and cost-effective approach to assessment of pituitary adrenocorticotropin and growth hormone reserve: combined use of the overnight metyrapone test and insulin-like growth factor-I standard deviation scores. *J Clin Endocrinol Metab* 2008;93:3763-8.
4. de Lind van Wijngaarden RF, Otten BJ, Festen DA, et al. High prevalence of central adrenal insufficiency in patients with Prader-Willi syndrome. *J Clin Endocrinol Metab* 2008;93:1649-54.